



Mycotoxin FAQs

1. What are mycotoxins?

Mycotoxins are natural chemicals produced by certain fungi, some of which cause ear rots in corn.

Mycotoxins are nonliving compounds that are byproducts that the fungi produce. Mycotoxins can have detrimental health effects to both humans and animals if they eat contaminated food or feed.




Figure 1. (A) Corn contaminated with *Aspergillus flavus*, the fungus responsible for Aspergillus ear rot. (B) The chemical structure of the mycotoxin aflatoxin.

2. What mycotoxins occur in corn?

There are five major mycotoxins associated with ear rot diseases of corn.

Aflatoxins are found in corn with *Aspergillus* ear rot (Figure 1).

Deoxynivalenol (DON, sometimes called vomitoxin) and **zearalenone** are found in corn with *Gibberella* ear rot (Figure 2).



Figure 2. Corn infected with the fungus that causes *Gibberella* ear rot.

Fumonisins are found in corn with *Fusarium* ear rot (Figure 3).

Ochratoxin is found in corn infected with *Penicillium verrucosum*, but some *Aspergillus* species also produce this mycotoxin.

3. What are the effects of mycotoxins?

The toxic effects of mycotoxins vary by type, dose, and animal species consuming the toxin.

Aflatoxin can affect the liver (hepatotoxicity), lead to cancer, and suppress the immune system.

Fumonisin can affect the liver (hepatotoxicity), lead to cancer, cause pulmonary edema (fluid in the lungs), and cause leukoencephalomalacia (irreversible, fatal brain damage) in horses and rabbits.




Figure 3. The chemical structure of a fumonisin molecule.

DON affects the gastrointestinal tract (which often makes animals vomit and refuse to feed) and can inflame the central nervous system.

Zearalenone can cause hyperestrogenism (estrogen overload), a condition that is particularly dangerous for female breeding animals.

Ochratoxin can cause cancer and is toxic to the kidneys.

4. What levels of mycotoxins are safe in food and feed?

The United States Food and Drug Administration (FDA) and Health Canada have issued **action levels** (legally allowed amounts) for aflatoxin and **advisory levels** (cautionary levels) for fumonisin and DON. The concentrations vary by mycotoxin and the intended use of the final corn products.

If grain or end products contain mycotoxin concentrations that exceed an action level, then the law may require the grain or products to be destroyed. Acceptable mycotoxin concentrations are most often set at the point of sale and in contract specifications. Corn for export must meet the limits set by each country, which are often more stringent than the United States.

Specific information about action and advisory levels are available from CornMycotoxins.com and the Canadian Food Inspection Agency, www.inspection.gc.ca/animals/feeds/regulatory-guidance/rg-8/eng/1347383943203/1347384015909?chap=1.

5. When should I test for mycotoxins?

If you have a field that has any ear rot problem, test the grain for mycotoxins. Testing for aflatoxin is recommended for corn that is severely drought stressed (Figure 4).



Figure 4. This corn is drought stressed, which can make it more prone to infection by the fungus that causes *Aspergillus* ear rot.

More information about grain sampling and mycotoxin testing is available in *Corn Disease Management: Grain Sampling and Mycotoxin Testing* (CPN-2003), available from the Crop Protection Network, CropProtectionNetwork.org.

6. How do I scout for potential problems?

Scout Fields Before Harvest

Scout fields for ear rot diseases at kernel maturity (black layer) and just before harvest.

Pay close attention to areas where the crop may have been stressed, damaged, or exposed to extreme environmental conditions. These areas include hillsides where drought stress may be more severe (for *Fusarium* and *Aspergillus* ear rots) and low areas where moisture from fog or high-dew conditions prevail (for *Gibberella* ear rot).

When you scout, randomly select plants and pull back the husks to examine the ears (Figure 5). A quick method is to select 100 plants across the field (20 ears each from five different areas). If you find a diseased ear, examine another 10 ears from adjacent plants.



Figure 5. Pull back the husk on corn ears to look for signs and symptoms of corn ear rots.

During your examination, ask three questions:

1. What ear rot disease is present?
2. How much of the ear is affected by ear rot (what is the severity of the disease)?
3. What proportion of the crop is affected?

Once you answer these questions, determine your next steps (see below).

More information about identifying ears rots is available in *Corn Disease Management: Ear Rots* (CPN-2001), available from the Crop Protection Network, CropProtectionNetwork.org.

Know When to Test for Mycotoxins

The risk of mycotoxins in harvested grain increases with the number of infected kernels on an ear. You should assume that diseased kernels contain mycotoxins.

However, mycotoxin levels may vary among diseased ears. Additionally, corn that appears to have a mild ear rot may still have very high mycotoxin levels. This makes the decision to test for mycotoxins difficult.

The threshold for aflatoxin should be very low. If you find any ears with *Aspergillus* ear rot, test the harvested grain for aflatoxin.

For DON and fumonisins, the threshold is flexible. If 30 percent of the ears you examine in a field have *Gibberella* or *Fusarium* ear rots, test for DON and fumonisins. You should also test your corn if you observe severe symptoms (more than 50 percent of the ear covered with mold) of either disease on multiple ears.

7. What are the mycotoxin testing options?

Mycotoxins are complex chemical compounds, which makes them difficult to quantify. However, several technologies can assess mycotoxin concentrations in corn grain.

For this reason, you should never rely solely on visual methods to confirm the presence of mycotoxins. A common visual test — the black light test — can indicate the presence of the fungus *Aspergillus flavus*, but it does not detect the aflatoxin it produces (Figure 6).



Figure 6. When crushed corn grain (top) is exposed to a black light, it can fluoresce (glow) if kojic acid is present (bottom right). *A. flavus* often produces kojic acid, but a black light test is not a reliable method for detecting aflatoxin.

For an accurate assessment, send grain samples to a professional laboratory for analysis (Figure 7). Local laboratories and grain inspection services may test individual corn samples for mycotoxins; however, sample testing can be expensive. Check with your local extension personnel for a more complete list of grain testing facilities in your area. The cost and submission procedures will vary by provider.

See *Corn Disease Management: Grain Sampling and Mycotoxin Testing* (CPN-2003) for more information.



Figure 7. Sample corn to send out for laboratory testing.

8. Will drying, heating, freezing, or applying chemicals reduce mycotoxins in grain?

No. Mycotoxins are extremely stable and heating, freezing, roasting, or treating with chemicals cannot reliably reduce mycotoxin levels within kernels. In some cases, you can reduce the overall mycotoxin concentration in the grain by removing broken grain (fines), foreign materials, and lightweight moldy kernels.

You can greatly reduce the further accumulation of mycotoxins in harvested grain by properly drying corn to less than 15 percent moisture. Dry grain to 13 percent for long-term storage.

When grain moisture levels are greater than 16 percent, the risk of aflatoxin and ochratoxin accumulation increases. Moisture greater than 18 percent elevates the risk of DON, zearalenone, and fumonisins. Warm conditions will accelerate the rates of spoilage and mycotoxin accumulation. There is no evidence that mycotoxin levels will increase in grain stored at an appropriately low moisture (Figure 8).




Figure 8. Healthy corn in a grain wagon.

It's important to point out that while high-temperature drying will stop mold growth and mycotoxin production, it will not reduce the level of mycotoxins already present. Quick drying is preferred over low-heat drying. Be wary of low-temperature, in-bin dryers for moldy corn, and be sure to meet proper ventilation requirements for dry corn storage.

More information about proper grain storage is available in *Corn Disease Management: Storing Mycotoxin-affected Grain* (CPN-2004), available from the Crop Protection Network, CropProtectionNetwork.org.

Find Out More

The Crop Protection Network (CPN) is a multi-state and international collaboration of university and provincial extension specialists, and public and private professionals who provide unbiased, research-based information to farmers and agricultural personnel. Our goal is to communicate relevant information that will help professionals identify and manage field crop diseases.

Find more crop disease resources at CropProtectionNetwork.org.

Authors

Tom Allen, Mississippi State University
 Martin Chilvers, Michigan State University
 Travis Faske, University of Arkansas
 Anna Freije, Purdue University
 Tom Isakeit, Texas A&M University
 Daren Mueller, Iowa State University

Trey Price, LSU AgCenter
 Damon Smith, University of Wisconsin
 Albert Tenuta, OMAFRA
 Kiersten Wise, Purdue University
 Charles Woloshuk, Purdue University

Reviewers

Gary Bergstrom, Cornell University
 Alyssa Collins, Pennsylvania State University
 Andrew Friskop, North Dakota State University
 Doug Jardine, Kansas State University
 Hillary Mehl, Virginia Tech University
 Alison Robertson, Iowa State University

Photo Credits

All photos were provided by and are the property of the authors and contributors except Figure 4 by John Obermeyer, Purdue University; Figures 5 and 6A by Burt Bluhm, University of Arkansas; and Figure 8 by Adam Sisson, Iowa State University.

Acknowledgements

Funding for this project was provided by the United States Department of Agriculture National Institute for Food and Agriculture (USDA-NIFA) project Integrated Management Strategies for Aspergillus and Fusarium Ear Rots of Corn. NIFA Award Number: 2013-68004-20359. We also thank the Grain Farmers of Ontario for support.

Design and production by Purdue Agricultural Communication.



United States Department of Agriculture
 National Institute of Food and Agriculture





The information in this publication is only a guide, and the authors assume no liability for practices implemented based on this information. Reference to products in this publication is not intended to be an endorsement to the exclusion of others that may be similar. Individuals using such products assume responsibility for their use in accordance with current directions of the manufacturer.

July 2016

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

©2016 by the Crop Protection Network. All rights reserved.



The Crop Protection Network is a multi-state and international collaboration of university/provincial extension specialists and public/private professionals that provides unbiased, research-based information to farmers and agricultural personnel.

